

# Measuring the weak mixing angle with the P2 experiment at MESA

BERGER N.<sup>1</sup>, AULENBACHER K.<sup>1,2</sup>, BAUNACK S.<sup>1</sup>, BECKER D.<sup>1</sup>, DIEFENBACH J.<sup>1,2</sup>, GERICKE M.<sup>3</sup>, GERZ K.<sup>1</sup>, HERBERTZ R.<sup>1</sup>, KUMAR K.<sup>4</sup>, MAAS F.<sup>1,2</sup>, MOLITOR M.<sup>1</sup>, PIÑEIRO D. R.<sup>1,2</sup>, SOROKIN I.<sup>1</sup>, SOUDER P.<sup>5</sup>, SPIESBERGER H.<sup>6</sup>, TYUKIN A.<sup>1</sup>, TYUKIN V.<sup>1</sup>, ZIMMERMANN M.<sup>1</sup>

(1. PRISMA Cluster of Excellence and Institute of Nuclear Physics, Johannes Gutenberg University, Mainz, Germany;

2. Helmholtz Institut, Mainz, Germany; 3. Department of Physics and Astronomy, University of Manitoba, Winnipeg, Canada;

4. Department of Physics and Astronomy, Stony Brook University, Stony Brook, USA; 5. Physics Department, Syracuse University, Syracuse, USA;

6. PRISMA Cluster of Excellence and Institute of Physics, Johannes Gutenberg University, Mainz, Germany)

**Abstract:** The P2 experiment in Mainz aims to measure the weak mixing angle  $\sin^2\theta_w$  in electron-proton scattering to a precision of 0.13%. In order to suppress uncertainties due to proton structure and contributions from box graphs, both a low average momentum transfer  $Q^2$  of  $4.5 \times 10^{-3} \text{ GeV}^2/c^2$  and a low beam energy of 155 MeV are chosen. In order to collect the enormous statistics required for this measurement, the new Mainz energy recovery superconducting accelerator (MESA) is being constructed. These proceedings describe the motivation for the measurement, the experimental and accelerator challenges and plans to tackle them.

**Key words:** weak mixing angle; parity violation; electron scattering

**CLC number:** O572.3      **Document code:** A      doi:10.3969/j.issn.0253-2778.2016.06.006

**Citation:** BERGER N, AULENBACHER K, BAUNACK S, et al. Measuring the weak mixing angle with the P2 experiment at MESA[J]. Journal of University of Science and Technology of China, 2016,46(6): 481-487.

BERGER N, AULENBACHER K, BAUNACK S, 等. MESA P2 实验对弱混合角的测量[J]. 中国科学技术大学学报, 2016,46(6): 481-487.

## MESA P2 实验对弱混合角的测量

BERGER N.<sup>1</sup>, AULENBACHER K.<sup>1,2</sup>, BAUNACK S.<sup>1</sup>, BECKER D.<sup>1</sup>, DIEFENBACH J.<sup>1,2</sup>, GERICKE M.<sup>3</sup>, GERZ K.<sup>1</sup>, HERBERTZ R.<sup>1</sup>, KUMAR K.<sup>4</sup>, MAAS F.<sup>1,2</sup>, MOLITOR M.<sup>1</sup>, PIÑEIRO D. R.<sup>1,2</sup>, SOROKIN I.<sup>1</sup>, SOUDER P.<sup>5</sup>, SPIESBERGER H.<sup>6</sup>, TYUKIN A.<sup>1</sup>, TYUKIN V.<sup>1</sup>, ZIMMERMANN M.<sup>1</sup>

(1. 德国约翰内斯堡大学核物理学院卓越研究中心, 美因茨, 德国; 2. 亥姆霍兹公司, 美因茨, 德国;

3. 曼尼托巴大学天文物理学系, 温尼伯格 加拿大; 4. 石溪大学天文物理学系, 石溪, 美国;

5. 锡拉丘兹大学物理系, 锡拉丘兹, 美国; 6. 德国约翰内斯堡大学物理学院卓越研究中心, 美因茨, 德国)

**Received:** 2015-11-30; **Revised:** 2016-04-20

**Foundation item:** Supported the Cluster of Excellence PRISMA and the Collaborative Research Center 1044, both Funded Through the Deutsche Forschungsgemeinschaft (DFG).

**Biography:** BERGER N (corresponding author), Professor/PhD. Research field: high energy physics. E-mail: niberger@uni-mainz.de

**摘要:**美因茨的 P2 实验致力于通过电子-质子散射将弱混合角  $\sin^2\theta_W$  的测量精确度提高到 0.13%。为了压低由于质子结构和箱图贡献的不确定性,选择了低平均四动量转移  $Q^2$  ( $4.5 \times 10^{-3} \text{ GeV}^2/c^2$ ) 和低束流能量 (155 MeV)。为了达到实验要求的巨大统计量,新的美因茨能量恢复超导加速器 (MESA) 正在建设中。本文描述了测量的动机、实验和加速器面临的挑战以及解决问题的方案。


**关键词:**弱混合角; 宇称破缺; 电子散射

## 0 Introduction

The weak mixing angle  $\sin^2\theta_W$  is one of the fundamental parameters of the standard model (SM) of elementary particle physics. It has been measured with great precision at the  $Z$  resonance<sup>[1]</sup>, where the determinations from LEP and SLD are marginally consistent. Due to quantum corrections, the effective weak mixing angle is a scale dependent quantity and measurements at different scales become important both for testing the SM and searching for effects of new physics beyond the SM in the running<sup>[2-3]</sup>.

Measurements at lower scales were obtained in neutrino nucleon scattering<sup>[4]</sup>, deep inelastic electron scattering<sup>[5]</sup>, parity violating electron scattering on electrons<sup>[6]</sup> and protons<sup>[7]</sup> and atomic parity violation in Caesium<sup>[8]</sup>. These measurements were sufficient to establish the running of  $\sin^2\theta_W$ , more precision is however required for a stringent test of the SM and searches for new physics. The final result from the  $Q_{\text{weak}}$  experiment is eagerly awaited and should improve on the published result<sup>[7]</sup> by a factor of three to four. More precise determinations require new experimental approaches, such as the proposed Møller<sup>[9]</sup> and deep inelastic (SOLID<sup>[10]</sup>) scattering experiments at JLAB and the P2 experiment in Mainz, which will be described in the following. An overview of current and planned experiments together with the theory prediction<sup>[11]</sup> for the running of  $\sin^2\theta_W$  is shown in Fig. 1.

A precise determination of the weak mixing angle at low scales is sensitive to contributions of new physics beyond the SM which can change the running of  $\sin^2\theta_W$  via contributions of new gauge bosons, additional fermions, mixing terms<sup>[12]</sup> or the exchange of very heavy particles which can be parametrized as four-



**Fig. 1** Scale dependence of  $\sin^2\theta_W$  together with completed and planned experimental measurements

fermion contact interactions<sup>[2]</sup>. In the last case, P2 will be sensitive to scales up to 49 TeV, comparable to the experiments at the large hadron collider after collecting  $300 \text{ fb}^{-1}$  of integrated luminosity.


## 1 Requirements

The P2 experiment aims to determine  $\sin^2\theta_W$  with a precision of 0.13% by performing a measurement of the parity violating asymmetry  $A_{PV}$  in electron-proton scattering. This asymmetry between the cross-sections for left- and righthanded electrons  $\sigma_L$  and  $\sigma_R$  is determined by the weak charge of the proton  $Q_W$ :

$$A_{PV} = \frac{\sigma_L - \sigma_R}{\sigma_L + \sigma_R} = \frac{G_F Q^2}{4\sqrt{2}\alpha} (Q_W + F(Q^2)) \quad (1)$$

where  $G_F$  is the Fermi constant, the fine structure constant and  $Q^2$  the squared four-moment transfer, setting the scale. Contributions stemming from the fact that the proton is not a point-like particle are collected in  $F(Q^2)$  and are small for low values of  $Q^2$ , thus motivating an experiment at low momentum transfer. Further hadronic uncertainties come from box graphs such as the one shown in Fig. 2. These uncertainties with a weak dependence on the momentum transfer do however, increase steeply with rising center-of-mass

energy<sup>[13-14]</sup>, favouring a low beam energy.



**Fig. 2**  $\gamma$ -Z box graph for electron-proton scattering

The weak mixing angle is related to  $Q_w$  via

$$Q_w = 1 - 4 \sin^2 \theta_w \quad (2)$$

which implies by propagation of uncertainty that a 0.13% measurement of  $\sin^2 \theta_w$  requires a 1.5% measurement of  $Q_w$ , which also corresponds to the target uncertainty in the asymmetry. Due to the small weak charge of the proton and the small  $Q^2$ , the expected asymmetry is only 33 ppb, thus requiring a measurement with 0.44 ppb precision. The statistical uncertainty scales with the number of scattered electrons as  $\frac{1}{\sqrt{N}}$ , which in turn requires the observation of  $\mathcal{O}(10^{18})$  electrons. For sociological reasons, the total measurement time is limited to 10 000 hours, which requires observing  $\mathcal{O}(10^{11})$  signal electrons per second.

These very high rates can be achieved by directing a 150  $\mu\text{A}$  electron beam onto a 60 cm long liquid hydrogen target, producing a luminosity of  $2.4 \times 10^{39} \text{ s}^{-1} \text{ cm}^{-2}$ .

The aim of determining  $\sin^2 \theta_w$  with a precision of 0.13% is thus extremely challenging for the accelerator and detector systems. The following sections outline how the MESA accelerator, the polarisation measurement and the P2 experiment intend to tackle these challenges.

## 2 The MESA accelerator

In order to accommodate the very long running time and demanding stability requirements of the P2 experiment, a new accelerator, the Mainz energy-recovery superconducting accelerator (MESA<sup>[15]</sup>) is being built.

With a maximum extracted beam energy of 155

MeV, MESA is small enough to fit into the existing halls that have become available with the completion of the A4 parity violating electron scattering program at the Mainz Microtron MAMI. P2 and the MAGIX spectrometer (see the contribution of A. Denig to this conference for details on the MESA program beyond the P2 experiment) will be housed in a new hall as part of the recently funded centre for fundamental physics. Fig. 3 shows the overall layout of accelerator and experiments (indicating existing and new halls at the institute of nuclear physics in Mainz).

P2 requires a highly polarized ( $> 85\%$ ), high intensity (150  $\mu\text{A}$ ) beam of 155 MeV electrons with excellent availability ( $> 4000 \text{ h/year}$ ). The beam helicity will be flipped several thousand times a second. The main challenge is to reduce any helicity correlated changes in beam intensity, energy, position and angle to less than 0.1 ppb. Here we can profit from the extensive experience in beam stabilization gained at the Mainz Microtron MAMI. Tab. 1 compares the values for helicity correlated beam fluctuations achieved at MAMI with the requirements for P2 at MESA. Whilst the energy stability already fulfills the demands, improvements of one to two orders of magnitude have to be achieved for position, angle and intensity; new digital feedback electronics for beam stabilization are currently being designed and tested at MAMI.

**Tab. 1** Helicity correlated beam fluctuations

Beam Quantity	Achieved at MAMI	Contribution to $\delta(A_{PV})$	Required for MESA
Energy	0.04 eV	$< 0.1 \text{ ppb}$	fulfilled
Position	3 nm	5 ppb	0.13 nm
Angle	0.5 rad	3 ppb	0.06 mrad
Intensity	14 ppb	4 ppb	0.36 ppb

The MESA lattice design is finalized, the superconducting RF cavities have been ordered and civil construction on the new hall will start 2016. We plan to start installing the accelerator in 2018 and have beam available for P2 before 2020.

## 3 Polarimetry

P2 requires a knowledge of the beam polarisation




Fig. 3 Layout of the MESA accelerator and experiments

of better than 0.5%. We aim to achieve this precision via two paths<sup>[16]</sup>, namely an invasive double Mott polarimeter at the electron source and a hydro-Møller polarimeter, which can be operated at the same time as and is placed right in front of the main experiment.

### 3.1 Double Mott polarimeter

The asymmetry of Mott scattering in thin foils can be used to determine the beam polarization, it however requires a precise knowledge of the analyzing power of the scattering foils which introduces a large uncertainty into the measurement. Double Mott scattering<sup>[17]</sup> in two foils allows to determine the effective analyzing power within the setup, thus reducing the associated uncertainty which makes it a suitable choice for precise source polarimetry at MESA. A prototype of the double-Mott polarimeter is currently tested with the MESA source prototype in operation in Mainz.

### 3.2 Hydro-Møller polarimeter

We plan to determine the beam polarization at the final energy with a hydro-Møller polarimeter<sup>[18]</sup> right in front of the main experiment. Here the asymmetry in Møller scattering of the beam electrons with the electrons in fully polarized atomic hydrogen is used. The hydrogen is polarized using a 7-8 T solenoid magnet. In order to avoid hydrogen recombination, the gas is kept at cryogenic temperatures and the walls of the vessel are coated with superfluid helium. Operating this cryogenic setup with a high intensity electron beam passing through the center is certainly challenging. The

cryostat/magnet for this setup is currently under construction.

## 4 The P2 experiment

For a given electron beam energy, the scattering angle  $\theta$  determines the momentum transfer  $Q^2$ . At low  $Q^2$ , the uncertainty of the asymmetry measurement is dominated by statistics and helicity correlated beam fluctuations. At large  $Q^2$ , uncertainties in the proton form factors become dominant. For our setup, the best accuracy can be reached with a central scattering angle of  $35^\circ$  at an angular acceptance of  $20^\circ$  (as shown in Fig. 4).




Fig. 4 Contributions to the uncertainty

In Fig. 4, contributions to the uncertainty of the  $\sin^2 \theta_W$  measurement at fixed beam energy, intensity and run time in dependence of the central scattering angle.  $G_E^S$ ,  $G_M^S$  and  $G_P^A$  refer to the uncertainties stemming from the electric and magnetic strange form

factor and the axial form factor of the proton respectively.

As the very high intensity beam produces several thousand Bremsstrahlung photons for every electron scattered into the angular range of interest as well as a large amount of Møller scattered electrons with low transverse momentum, a magnetic spectrometer is required in order to guide signal electrons to detectors whilst at the same time shielding them from photon and Møller backgrounds. The detectors in turn face the challenge of reliably detecting more than 100 GHz of scattered electrons. The following sections will describe the spectrometer design, the development of integrating Cherenkov detectors as well as a pixellated tracking detector for a precise determination of the momentum transfer  $Q^2$ .

#### 4.1 Spectrometer

For the spectrometer design, the main choice is between a toroidal (as used in the  $Q_{\text{weak}}$  experiment) and a solenoidal magnetic field. The advantages of a toroidal setup, such as zero field in the target region and easy access to instrumentation are compromised by the fact that the coils are necessarily inside of the spectrometer acceptance, typically leading to a loss of about half the signal electrons and consequently a doubling of the measurement time, which is unrealistic in the context of P2.

We have thus decided to employ a solenoidal design and studied possible placements of the target, shielding and detectors for several existing solenoids, e. g. from the ZEUS experiment at HERA<sup>[19]</sup> or the FOPI experiment at GSI<sup>[20]</sup>. We have shown using both ray-tracing in the magnet field maps and full Geant4 based simulations that with a careful optimization of the shape and placement of lead shields, sufficient signal-to-background ratios can be achieved in the integrating detectors. A possible view of the setup is shown in the rendering in Fig. 5 (showing the solenoid coil, target, lead shielding and integrating detectors as well as a hunky physicist for scale).

#### 4.2 Integrating detectors

Individually counting hundreds of GHz of




Fig. 5 Rendering of a possible P2 setup

electrons is extremely challenging, but not actually required for P2. Instead, we opt for an integrating measurement, where the electrons produce Cherenkov light in bars of fused silica (quartz), which is detected with photomultipliers operated at low gain. The current of these photomultipliers is integrated over one helicity period and read out with high precision (22 bit) analog-to-digital converters.

We are currently testing different types and polishing finishes as well as wrappings of quartz bars with the MAMI beam and have found both a performance sufficient for P2 as well as an excellent agreement of the light yield at different incident angles with Geant4 based simulations.

The switchable gain photomultiplier base is currently under development in Mainz; for the integrating ADC a joint development with the Møller experiment is ongoing at the University of Manitoba.

#### 4.3 Tracking detectors

In order to determine the average momentum transfer  $\langle Q^2 \rangle$  of the scattered electrons creating signals in the integrating detectors, a tracking detector is required. The high rates at MESA, the precision requirements and the low momenta of the scattered electrons (making multiple coulomb scattering in the tracker material the dominating uncertainty in the momentum measurement) call for a fast, high granularity sensor with very little material. We choose to employ high-voltage monolithic active pixel sensors (HV-MAPS<sup>[21-25]</sup>) as the detector technology. These sensors, manufactured in a commercial CMOS

technology, apply a “high” voltage of around 90 V between deep  $n$ -wells and the substrate, leading to very fast charge collection from a thin depletion layer. The thin charge collection zone allows for thinning of the sensors to just 50  $\mu\text{m}$ . The sensor is segmented into 80 by 80  $\mu\text{m}$  pixels. The CMOS process used allows for integrating both analog and digital electronics directly on the sensor; the output are zero suppressed hit addresses and timestamps on a fast differential link. In the development of the sensors for P2, we closely collaborate with the Mu3e, ATLAS and Panda experiments.

From these thin sensors, we plan to build a tracking detector with four planes (shown in Fig. 6, consisting of four planes of HV-MAPS sensors). The arrangement in two double planes combines a good momentum and angular resolution in a multiple scattering dominated regime with ease of reconstruction in a high multiplicity environment. We are currently studying tracking algorithms that also perform well in the non-uniform field close to the edge of the magnet and are at the same time robust and fast enough to allow for on-line track finding and fitting, possibly on highly parallel architectures such as graphics processing units (GPUs).




Fig. 6 Schematic view of the P2 tracking detector

## 5 Conclusion

The P2 experiment aims to measure  $\sin^2\theta_W$  at low momentum transfer with unprecedented accuracy, which both improves the precision on one of the fundamental parameters of the standard model and allows to search for new physics. The new MESA accelerator in Mainz will provide a stable, very high intensity electron beam combined with precision polarimetry. P2 will measure the parity violating asymmetry in electron-proton scattering using a

solenoid spectrometer with integrating Cherenkov detectors combined with a thin pixel tracker. Accelerator commissioning is scheduled to start in 2018 and a first P2 data taking for 2020. Beyond the  $\sin^2\theta_W$  measurement, P2 can also be used to study parity violation with different target materials, giving access e. g. to neutron skins. The MESA accelerator also has a wide physics program beyond P2, described elsewhere in these proceedings.

## References

- [1] SCHAEEL S, BARATE R, BRUNELIERE R, et al. Precision electroweak measurements on the Z resonance [J]. *Physics Reports*, 2006, 427(5-6): 257-454.
- [2] ERLER J, KURYLOV A, RAMSEY-MUSOLF M J. The weak charge of the proton and new physics [J]. *Physical Review D*, 2003, 68(1): 462-467.
- [3] ERLER J, RAMSEY-MUSOLF M J. Low energy tests of the weak interaction [J]. *Progress in Particle Nuclear Physics*, 2004, 54(2): 351-442.
- [4] ZELLER G P, MCFARLAND K S, ALTON A, et al. Precise determination of electroweak parameters in neutrino-nucleon scattering [J]. *Physical Review Letters*, 2001, 88(9): 386-389.
- [5] AKTAS A, ANDREEV V, ANTHONIS T, et al. A determination of electroweak parameters at HERA [J]. *Physics Letters B*, 2006, 632(1): 35-42.
- [6] ANTHONY P L, ARNOLD R G, ARROYO C, et al. Observation of parity nonconservation in mller scattering [J]. *Physical Review Letters*, 2004, 92(18): 181602.
- [7] ANDROIC D. First determination of the weak charge of the proton [J]. *Physical Review Letters*, 2013, 111(14): arXiv:1307.5275.
- [8] WOOD C S, BENNETT S C, CHO D, et al. Measurement of parity nonconservation and an anapole moment in cesium [J]. *Science*, 1997, 275953070: 1759-1763.
- [9] BENESCH P, BRINDZA R D, CARLINI J P, et al. The MOLLER experiment: An ultra-precise measurement of the weak mixing angle using Mller scattering [J]. *EPrint Arxiv*, 2014; arXiv:1411.4088.
- [10] CHEN J P, GAO H, HEMMICK T K, et al. A white paper on SoLID (Solenoidal large intensity device) [J]. *EPrint Arxiv*, 2014; arXiv:1409.7741.
- [11] ERLER J, RAMSEY-MUSOLF J. Weak mixing angle at low energies [J]. *Physical Review D*, 2004, 72(7): arXiv:hep-ph/0409169.
- [12] DAVOUDI ASL H, LEE H S, MARCIANO W J. Muon

- $g-2$ , rare kaon decays, and parity violation from dark Bosons[J]. *Physical Review D*, 2014, 89(9):98-105.
- [13] GORCHTEIN M, HOROWITZ C J, RAMSEY-MUSOLF M J. Model-dependence of the  $\gamma Z$  dispersion correction to the parity-violating asymmetry in elastic  $ep$  scattering[J]. *Physical Review C* 84, 2011, 84(1): 015502 (arXiv: 1102.3910).
- [14] GORCHTEIN M, SPIESBERGER H, ZHANG X. How strange is pion electroproduction? [J]. *Physical Letters B*, 2016, 752: 135-145.
- [15] AULENBACHER K. The MESA accelerator [C]// AIP Conference Proceedings. Cambridge, USA: 2013, 1563: 5-12.
- [16] AULENBACHER K, ALEXANDER I, TIOUKINE V. The polarimetry chain for the P2 experiment [J]. *Nuovo Cimento C*, 2012, 186: 035N04.
- [17] GELLERICH A, KESSLER J. Precision measurement of the Sherman asymmetry function for electron scattering from gold [J]. *Physical Review A*, 1991, 43(1): 204-216.
- [18] BARTOLOMÉ P A, AULENBACHER K, TYUKIN V. Møller polarimetry with polarized atomic hydrogen at MESA [C]// AIP Conference Proceedings. 1553, 243-246.
- [19] ACERBI E, ALESSANDRIA F, BACCAGLIONI G, et al. The Zeus thin superconducting solenoid [J]. *Proceedings of International Conference on Magnet Technology*. Zürich, Switzerland, 1985: 163-166.
- [20] RITMAN J, HERMANN N, BEST D, et al. On the transverse momentum distribution of strange hadrons produced in relativistic heavy ion collisions [J]. *Nuclear Physics Proceedings*, 1995, 44(S): 708-715.
- [21] PERIĆ I. A novel monolithic pixelated particle detector implemented in high-voltage CMOS technology [J]. *Nuclear Instruments and Methods in Physics Research Section A*, 2007, 582: 876-885.
- [22] PERIĆ I, FISCHER P, KREIDL C, et al. High-voltage pixel detectors in commercial CMOS technologies for ATLAS, CLIC and Mu3e experiments [J]. *Nuclear Instruments and Methods in Physics Research Section A*, 2013, 731(18): 131-136.
- [23] SHRESTHA S (Mu3e Collaboration). The high-voltage monolithic active pixel sensor for the Mu3e experiment [C]// *Proceedings of Technology and Instrumentation in Particle Physics*. Amsterdam, 2014: 047(1-9).
- [24] AUGUSTIN H, BERGER N, BRAVAR S, et al. The MuPix high voltage monolithic active pixel sensor for the Mu3e experiment [J]. *Journal of Instrumentation*, 2015, 10(3): C03044(1-8).
- [25] PERIĆ I, EBER R, EHRLER F, et al. Overview of HVCMOS pixel sensors [J]. *Journal of Instrumentation*, 2015, 10(5): C05021(1-8).