



Nickel-catalyzed alkene *ipso*-selective reductive hydroamination with nitroarenes


Zhen Li, Jiawang Wang, Xi Lu , and Yao Fu 

Department of Applied Chemistry, University of Science and Technology of China, Hefei 230026, China

Correspondence: Xi Lu, E-mail: luxl@mail.ustc.edu.cn; Yao Fu, E-mail: fuyao@ustc.edu.cn

© 2022 The Author(s). This is an open access article under the CC BY-NC-ND 4.0 license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Graphical abstract





Aromatic amines with primary and secondary alkyl groups could be accessed through nickel-catalyzed *ipso*-selective alkene hydroamination with nitroarenes.

Public summary

- A nickel-catalyzed *ipso*-selective alkene hydroamination with nitroarenes under mild reductive conditions was developed.
- Repaid preparation of aromatic amines with primary and secondary alkyl groups and compatibility with many functional groups were demonstrated.
- Based on the literature and mechanistic investigation, a plausible reaction mechanism involving in situ generation of alkylnickel intermediates and reaction with in situ generated nitrosoarenes was proposed.

Nickel-catalyzed alkene *ipso*-selective reductive hydroamination with nitroarenes

Zhen Li, Jiawang Wang, Xi Lu , and Yao Fu 

Department of Applied Chemistry, University of Science and Technology of China, Hefei 230026, China

 Correspondence: Xi Lu, E-mail: luxl@mail.ustc.edu.cn; Yao Fu, E-mail: fuyao@ustc.edu.cn

© 2022 The Author(s). This is an open access article under the CC BY-NC-ND 4.0 license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).



Cite This: *JUSTC*, 2022, 52(12): 1 (9pp)



Read Online



Supporting Information

Abstract: Aromatic amine synthesis via reductive coupling between alkenes and nitroarenes is attractive; however, it remains underdeveloped. Herein, we report a nickel-catalyzed alkene hydroamination with nitroarenes under mild reductive conditions. This reaction exhibited an *ipso*-selectivity and enabled rapid preparation of aromatic amines with primary and secondary alkyl groups. Many functional groups were well tolerated, providing an efficient approach for drug-like arylamine synthesis.

Keywords: nickel; alkene; nitroarene; reductive hydroamination; C-N coupling

CLC number: TQ246.3

Document code: A

1 Introduction

Aromatic amines are necessary organic chemicals in many fields, such as the chemical industry, materials, medicines, and agricultural and veterinary medicines^[1]. For example, the small-molecule pharmaceuticals palbociclib, apixaban, elixacator, and tofacitinib were among the top 200 drugs by retail sales in 2021 (Fig. 1a). Classical methods for synthesizing aryl or heteroaryl amines include amine-carbonyl reductive amination^[2], nucleophilic substitutions of amines with alkyl halides^[3–5], Buchwald–Hartwig C–N coupling reactions^[6–8], and Ullmann–Ma-type amination reactions^[9–14] (Fig. 1b). The methods mentioned above for synthesizing aromatic amines via C–N bond formation require primary aromatic amines as starting materials. However, primary aromatic amines are usually prepared by nitroarene derivative hydrogenation reduction^[15,16]. Therefore, the direct synthesis of aromatic amines using large industrial nitroarenes shortens the synthesis steps and improves the synthetic efficiency^[16–20]. On the other hand, the stable features and extensive sources of alkenes^[21–28] have promoted alkene hydroamination as an attractive alternative for amine synthesis^[29–35]. In 2015, Baran et al.^[36] reported a practical arylamine synthesis via alkene hydroamination with nitroarenes, which combines iron-promoted hydrogen atom transfer, radical addition, and zinc reduction. In 2016, Hu et al.^[15] realized an elegant single-step (hetero)aryl amine synthesis via iron-catalyzed reductive coupling between alkyl halides and nitroarenes. In 2018, Zhu et al.^[37] developed an efficient benzylic arylamine synthesis via sequential nickel-catalyzed chain-walking and hydroamination processes. In 2021, Wang and Zhu et al.^[38] extended this efficient alkene hydroamination reaction to an asymmetric version. Very recently, Wang and Zhu et al.^[39] developed a useful method to access α -aminoboronates via hydroamidation of alkenyl boronates. Despite these achievements in synthesiz-

ing aromatic amines with tertiary or benzylic alkyl groups (Fig. 1c), general methods for synthesizing aromatic amines with primary and secondary alkyl groups from nitroarenes with alkenes remain underdeveloped.

Our group systematically researched base-metal-catalyzed alkene functionalization reactions, catalytic mechanistic studies and applications in functional organic molecule synthesis^[40–44]. Based on our research interests in alkene hydroalkylation reactions (Fig. 1d)^[45–51], we envisioned that nickel-promoted reductive hydroamination of alkenes and nitroarenes might be suitable for achieving our amine synthesis goal. Herein, we report a nickel-catalyzed alkene *ipso*-selective reductive hydroamination with nitroarenes to access aromatic amines with primary and secondary alkyl groups (Fig. 1e).

2 Materials and methods


2.1 Materials

The following chemicals were purchased and used as received: nickel(II) bromide 2-methoxyethyl ether complex (CAS: 312696-09-6, Aldrich, 459674-5G); triethoxysilane (CAS: 998-30-1, TCI, T1040); diethoxymethylsilane (CAS: 27176-10-9, Adamas, 35415E); tetrahydrofuran (CAS: 109-99-9, Energy Chemical, W3100755000); 1,4-dioxane (CAS: 123-91-1, Adamas, 01375906); methanol (CAS: 67-56-1, Sinopharm, 10014118); *N,N*-dimethylacetamide (CAS: 127-19-5, Adamas, 011342855); potassium fluoride (CAS: 7789-23-3, Acros, 01163384); and sodium carbonate anhydrous (CAS: 497-19-8, Sinopharm, 10019260).


2.2 Analytical methods

¹H NMR, ¹³C NMR, ¹¹B NMR and ¹⁹F NMR spectra were recorded on a Bruker 400 MHz spectrometer and a Bruker 500 MHz spectrometer at 295 K in CDCl₃ unless otherwise noted.


(a) representative drug molecules containing (hetero) aromatic amine structure




(b) classical methods for synthesizing aryl or heteroaryl amines




(c) state of the art: direct synthesis of (hetero) aromatic amines using nitro(hetero)arenes



(d) our previous works: nickel- and cobalt-catalyzed alkene (alkyne) hydroalkylation



(e) this work: alkene *ipso*-selective reductive hydroamination with nitroarenes



- introducing alkyl groups with alkenes
- general method for primary and secondary alkyl groups
- broad substrate scope of nitroarenes and alkenes
- introducing aniline structures with nitroarenes
- moderate to good yields and regioselectivities
- general method for aromatic amine synthesis

Fig. 1. State-of-the-art aromatic amine synthesis and our strategy. DG = directing group; FG = functional group; DEMS = diethoxymethylsilane; diglyme = 2-methoxyethyl ether; TBAI = tetrabutylammonium iodide.

Data for ¹H NMR were reported as follows: chemical shift (δ ppm), multiplicity, coupling constant (Hz), and integration. Data for ¹³C NMR were reported as follows: chemical shift (δ ppm), multiplicity, and coupling constant (Hz). Data for ¹¹B NMR were reported as follows: chemical shift (δ ppm), multiplicity, and coupling constant (Hz). Data for ¹⁹F NMR were

reported as follows: chemical shift (δ ppm), multiplicity, coupling constant (Hz). Chemical shifts were reported using the residual solvent CHCl₃ as the internal reference for ¹H NMR (δ = 7.260 ppm) and the CDCl₃ peak as the internal reference for ¹³C NMR (δ = 77.160 ppm). High-resolution mass spectral analysis (HRMS) data were acquired on a Water

XEVO G2 Q-TOF (Waters Corporation). Gas chromatographic (GC) analysis was acquired on a Shimadzu GC-2010 plus Series GC system equipped with a flame-ionization detector. Organic solutions were concentrated under reduced pressure on a Büchi rotary evaporator. Column chromatographic purification of the products was accomplished using forced-flow chromatography on silica gel (300–400 mesh).

2.3 General procedure

In air, a 10 mL screw-cap test tube equipped with a magnetic stirrer was charged with **L8** (0.03 mmol, 15 mol%), NiBr₂(diglyme) (0.02 mmol, 10 mol%), Na₂CO₃ (0.6 mmol, 3.0 equiv.), TBAI (0.1 mmol, 0.5 equiv.), and Mg powder (0.1 mmol, 0.5 equiv.) (if the nitroarene or the alkene was solid, they were added at this step). The test tube was evacuated and backfilled with argon three times. Then, solvent (1,4-dioxane : MeOH = 5 : 1, 1.20 mL) was added, followed by alkene (0.4 mmol, 2.0 equiv.) and nitroarene (0.2 mmol, 1.0 equiv.). Then, DEMS (2.0 mmol, 10.0 equiv.) was added dropwise via a syringe, and the solution was stirred for 5 min at 25 °C, followed by stirring at 50 °C for 12 h. The reaction mixture was diluted with H₂O followed by extraction with EtOAc, dried with anhydrous Na₂SO₄ and concentrated in vacuo. The residue was purified by flash column chromatography on silica gel to obtain the target product. The yield represents the isolated yield of the major product. The reaction regioselectivity was determined by GC analysis of the reaction mixture. In a few cases, the reaction regioselectivity could not be determined by GC analysis; ¹H NMR determined regioisomeric ratios after chromatography purification.



3 Results and discussion

3.1 Screening of reaction conditions

As shown in Table 1, we began this study with the synthesis of *N*-hexylaniline (**3**) from nitrobenzene (**1**) and hex-1-ene (**2**) using the hydroalkylation reaction conditions reported in our early works^[45]. With a merely increased amount of silane (entry 1), we obtained a mixture of linear product **3** and branched product **4** in a 39% GC (gas chromatography) yield and a 3.4 : 1.0 regioselectivity (*l* : *b*, linear : branched). Then, we screened the influence of all reaction parameters, such as nickel catalysts, ligands, silanes as hydride sources, bases, additives, and solvents. Other nitrogen- or phosphine-containing ligands were examined: bipyridine (**L2**, entry 2), bioxazoline (**L4**, entry 4), and terpyridine (**L5**, entry 5) yielded only trace amounts of cross-coupling products; the ethylenediamine-structured ligand (**L3**, entry 3) provided a better 5.0 : 1.0 regioselectivity with a decreased total yield; and the phosphine ligand 1,2-bis(diphenylphosphino)ethane (dppe, entry 6) could be used with a decreased 29% GC yield and 1.9 : 1.0 regioselectivity. In solvent screening (entries 7–11), a low polarity solvent (entry 8) led to improved coupling efficiency; a 1,4-dioxane/MeOH mixed solvent further improved the yield (entry 11). The selection of NiBr₂(diglyme), Na₂CO₃, and diethoxymethylsilane (DEMS) was critical to this reaction: other nickel catalysts, bases, and silanes were much less effective (entries 12–18). Next, we tried to add additives to the reaction: the addition of tetrabutylammonium

iodide (TBAI, entries 19 and 20) could significantly improve the coupling efficiency; additional zinc (entry 21) or magnesium

Table 1. Optimization of reaction conditions.

entry	variation from standard conditions	yield/% ^a	3 : 4 (<i>l</i> : <i>b</i>) ^b
1	none	39	3.4 : 1.0
2	L2 instead of L1	<2	—
3	L3 instead of L1	12	5.0 : 1.0
4	L4 instead of L1	<2	—
5	L5 instead of L1	<2	—
6	dppe instead of L1	29	1.9 : 1.0
7	CH ₃ CN instead of DMAc	24	1.4 : 1.0
8	toluene instead of DMAc	50	2.8 : 1.0
9	THF instead of DMAc	25	2.5 : 1.0
10	1,4-dioxane instead of DMAc	45	3.6 : 1.0
11	1,4-dioxane : MeOH = 5 : 1 instead of DMAc	57	3.4 : 1.0
12	NiI ₂ instead of NiBr ₂ (diglyme)	49 ^c	2.5 : 1.0
13	Ni(acac) ₂ instead of NiBr ₂ (diglyme)	8 ^c	1.0 : 1.0
14	Ni(cod) ₂ instead of NiBr ₂ (diglyme)	18 ^c	1.0 : 3.5
15	KF instead of Na ₂ CO ₃	38 ^c	1.9 : 1.0
16	NaOH instead of Na ₂ CO ₃	<2 ^c	—
17	PMHS instead of DEMS	35 ^c	2.8 : 1.0
18	(MeO) ₃ SiH instead of DEMS	29 ^c	2.9 : 1.0
19	1.0 equiv. TBAI added	66 ^c	1.9 : 1.0
20	0.5 equiv. TBAI added	66 ^c	2.3 : 1.0
21	0.5 equiv. TBAI and 0.5 equiv. Zn powder added	68 ^c	4.2 : 1.0
22	0.5 equiv. TBAI and 0.5 equiv. Mg powder added	78 ^c	4.6 : 1.0
23	L6 instead of L1	38 ^d	6.7 : 1.0
24	L7 instead of L1	56 ^d	12 : 1.0
25	L8 instead of L1	85 (70) ^{d,e}	6.3 : 1.0
26	5.0 equiv. DEMS used	42	6.9 : 1.0
27	3.0 equiv. DEMS used	10	6.4 : 1.0

^aConditions: **1** (0.10 mmol, 1.0 equiv.), **2** (0.20 mmol, 2.0 equiv.), nickel catalyst (0.01 mmol, 10 mol%), ligand (0.015 mmol, 15 mol%), silane (1.0 mmol, 10.0 equiv.), base (0.30 mmol, 3.0 equiv.), additives, solvent (0.60 mL, 0.167 mol/L), 50 °C, 12 h. GC (gas chromatography) yield.

^bYields and regioisomeric ratios were determined by GC analysis with triphenylmethane as an internal standard. The yield refers to total yield for the mixture of the linear product (**3**) and branched product (**4**). ^c1,4-Dioxane : MeOH = 5 : 1 as solvent. ^d1,4-Dioxane : MeOH = 5 : 1 as solvent, 0.5 equiv. TBAI and 0.5 equiv. Mg powder were added. ^eIsolated yield for the linear product in parentheses. DEMS = diethoxymethylsilane; PMHS = polymethylhydrosiloxane; DMAc = *N,N*-dimethylacetamide; THF = tetrahydrofuran; acac = acetylacetonate; cod = 1,5-cyclooctadiene; dppe = 1,2-bis(diphenylphosphino)ethane; Bn = benzyl; TBAI = tetrabutylammonium iodide.

(entry 22) powder could simultaneously improve the coupling yield and regioselectivity. The iodine ion might have a protective effect on the nickel catalyst. Another alternative is the formation of alkyl iodides in the presence of iodine ions. Finally, we tried to fine-tune the ligand under the conditions of entry 22 to seek better results (entries 23–25). To our delight, the use of pyridine-oxazoline ligand (**L8**) delivered satisfactory results (entry 25): a combination of NiBr₂(diglyme) and **L8**, Na₂CO₃ as the base, DEMS as silane, TBAI and Mg

powder as additives, in a 1,4-dioxane/MeOH mixed solvent delivered linear product **3** with 85% GC yield for the linear and branched product, 70% isolated yield for the linear product, and good regioselectivity profile (6.3 : 1.0 *l* : *b*). If the DEMS loading could be reduced, it would greatly improve the synthetic utility of this methodology. Unfortunately, the low loadings (5.0 equiv. in entry 26 and 3.0 equiv. in entry 27) DEMS led to decreased coupling yields.




Fig. 2. Scope of substrates. Standard conditions: nitroarene (0.20 mmol, 1.0 equiv.), alkene (0.40 mmol, 2.0 equiv.), NiBr₂(diglyme) (0.02 mmol, 10 mol%), **L8** (0.03 mmol, 15 mol%), DEMS (2.0 mmol, 10.0 equiv.), Na₂CO₃ (0.60 mmol, 3.0 equiv.), TBAI (0.10 mmol, 0.5 equiv.), Mg powder (0.10 mmol, 0.5 equiv.), solvent (1,4-dioxane : MeOH = 5 : 1, 1.20 mL, 0.167 mol/L), 50 °C, 12 h. The yield represents the isolated yield of the major product. The reaction regioselectivity was determined by GC analysis of the reaction mixture. ^aFor these cases, the reaction regioselectivity could not be determined by GC analysis, ¹H NMR determined regioisomeric ratios shown in the table after chromatography purification. ^b**L1** instead of **L8** as ligand. DEMS = Diethoxymethylsilane; TBAI = Tetrabutylammonium iodide.

3.2 Scope of reductive hydroamination

With the optimized reaction conditions, we examined the substrate scope of the reductive hydroamination reaction (Fig. 2). In all cases, various alkenes and nitroarenes with different functional groups delivered the desired products in moderate to good yields. This reaction could be applied to terminal (3, 5–15) and internal (16–18) alkenes. Cyclic internal alkenes (16–18) performed well in this transformation; the ring size of cyclic alkenes did not significantly affect the coupling efficiency, even with extension ring substrates (18). Various nitroarenes (19–31) were suitable substrates and provided the

respective products with satisfactory yields. For the nitroarene coupling partners, electroneutral (3), electron-rich (19, 20, 23, and 27), and electron-deficient (24 and 25) substrates gave moderate to good yields. Under mild reaction conditions, many functional groups were well tolerated, such as ether (6 and 7), silyl ether (8), ester (9 and 24), keto carbonyl (10, 25, and 26), aryl fluoride (21), thioether (27), cyano (28), and even amide-possessing N–H bonds (11). Heterocycles such as phthalimide (12), furan (13), indole (14), and pyrrole (15) posed no problem. The compatibility to electrophilic aryl chloride (22), aryl bromide (23), aryl triflate




Fig. 3. Synthetic applications. Standard conditions are shown in Fig. 2. ^aThe yield represents the total GC yield. The reaction regioselectivity was determined by GC analysis of the reaction mixture. ^bThe yield represents the isolated yield of the major product. The reaction regioselectivity was determined by GC analysis of the reaction mixture. ^cYields reported in reference. ^dThe yield represents the isolated yield of the major product. Regioselectivity could not be determined by GC analysis. The regioisomeric ratio was determined to be >20 : 1.0 *l* : *b* by ¹H NMR after chromatography purification. DCMI-PHOS = dicyclohexyl(2-mesityl-1*H*-inden-1-yl)phosphine; DME = 1,2-dimethoxyethane; dba = dibenzylideneacetone.

(30), and nucleophilic arylboronate (29) further provided further functionalization possibilities via transition-metal-catalyzed cross-coupling reactions. In addition, this reaction could be conducted in the presence of an alcohol group (31) without protection. The good functional group compatibility and the simplicity of operation provided an efficient approach for drug-like arylamine synthesis.

3.3 Synthetic applications

We conducted the reaction between nitrobenzene (1) and (*E*)-hex-3-ene (2) as a model reaction to examine the reactivity and regioselectivity of acyclic internal alkenes (Fig. 3a). In this case, an isomer mixture was obtained, and linear product 3 was the major isomer. Our approach has a few advantages over former amine synthesis methods (Fig. 3b), such as *N*-alkylation of amines with alcohols via dehydration reaction (Reaction I)^[52], amine-carbonyl reductive amination (Reaction II)^[53], and Buchwald–Hartwig C–N coupling reactions (Reaction III)^[54]. To further demonstrate the synthetic utility, we evaluated the potential of this approach in the late-stage

modification of complex molecules (Fig. 3c). Alkenes derived from drug molecules or other biorelevant molecules, such as indomethacin (41), carprofen (42), and naproxen (43), underwent the reductive hydroamination process smoothly.

3.4 Mechanistic investigation

A series of control and mechanistic experiments were carried out for mechanistic investigation (Fig. 4). In the control experiments with catalysts (Fig. 4a), the model reaction could not occur without nickel catalysts or using cobalt or iron catalysts. We found that in the absence of an alkene, nitrobenzene (1) was reduced under the standard conditions (Fig. 4b). However, the reduction of nitrobenzene would not occur in the absence of nickel catalysts. In the control experiments with potential intermediates (Fig. 4c), a series of plausible intermediates, including aniline (34), nitrosobenzene (44), *N*-phenylhydroxylamine (45), azoxybenzene (46), and azobenzene (47), were subjected to standard conditions instead of nitrobenzene (1). Nitrosobenzene (44) delivered product 3 in 6% GC yield with a 1.1 : 1.0 regioselectivity. In addition, we




Fig. 4. Mechanistic investigation. Standard conditions: nitroarene or potential intermediate (0.20 mmol, 1.0 equiv. or 0.10 mmol, 0.5 equiv.), alkene (0.40 mmol, 2.0 equiv.), NiBr₂(diglyme) (0.02 mmol, 10 mol%), **L8** (0.03 mmol, 15 mol%), DEMS (2.0 mmol, 10.0 equiv.), Na₂CO₃ (0.60 mmol, 3.0 equiv.), TBAI (0.10 mmol, 0.5 equiv.), Mg powder (0.10 mmol, 0.5 equiv.), solvent (1,4-dioxane : MeOH = 5 : 1, 1.20 mL, 0.167 mol/L), 50 °C, 12 h. Yields and regioisomeric ratios were determined by GC analysis with triphenylmethane as an internal standard. Total yield for the mixture of the linear product and branched product.




Fig. 5. Plausible reaction mechanism.

conducted a control nitrosobenzene experiment with a slow dropping operation; the yield and regioselectivity of product **3** were improved. Combining the results obtained in Fig. 4c and related literature^[15,37], we suggested that the nitrosoarene was a viable reaction intermediate. On the other hand, alkene hydroamination with in situ-generated aniline was less likely because no desired product formed in the control experiment with aniline. We also conducted a control experiment to probe whether alkylsilane acted as an intermediate (Fig. 4d). Trimethoxy(octyl)silane (**48**) was treated with nitrobenzene (**1**) under standard conditions, and the desired product **5** was not observed. This result ruled out the possibility of in situ alkene hydrosilylation and C-N coupling between alkylsilanes and nitroarenes.

3.5 Plausible reaction mechanism

Based on the literature and our mechanistic investigation^[15,16,23,37,38,40,45–50], we proposed a plausible reaction mechanism, as outlined in Fig. 5. Active Ni(I)X species (A) could be generated from the initial Ni(II) catalysts. Ni(I)X species (A) delivered a nickel hydride species Ni(I)H (B) after the reaction with silane. Then, alkylnickel(I) intermediates (C) were generated through the regioselective migratory insertion of Ni(I)H across the alkene. Reaction of alkylnickel(I) intermediates (C) with the nitrosoarenes generated in situ occurred to access the hydroxylamine intermediates and regenerate the active Ni(I)X species (A)^[37]. Finally, the reduction of hydroxylamines gave aromatic amine products.

4 Conclusions

We developed a nickel-catalyzed alkene hydroamination with nitroarenes under mild reductive conditions. This reaction exhibited an *ipso*-selectivity and enabled the rapid preparation of aromatic amines with primary and secondary alkyl groups. Many functional groups were well tolerated, providing an ef-

ficient approach for drug-like arylamine synthesis under protecting-group-free conditions.

Supporting information

The supporting information for this article can be found online at <https://doi.org/10.52396/JUSTC-2022-0119>. The supporting information includes 65 figures. The characterization of all new compounds is provided in the supporting information.

Acknowledgements

This work was supported by the USTC Research Funds of the Double First-Class Initiative (YD3530002002), the Fundamental Research Funds for the Central Universities (WK2060000043 and WK3530000013), and the National Science Foundation of Anhui Province (2208085J26 and 2208085QB36). The authors thank Dr. Yuli He (China Pharmaceutical University) for insightful mechanistic discussion and Dr. Xiaoxu Wang (University of Science and Technology of China) for preliminary experiments.

Disclosures

Pay caution to eye protection when handling silanes, which might cause potential security risks. Silanes might be disproportionated and form pyrophoric SiH₄. Suitable precautions are needed.

Conflict of interest

The authors declare that they have no conflict of interest.

Biographies

Zhen Li is a master's student under the supervision of Prof. Yao Fu at the University of Science and Technology of China. His research mainly focuses on nickel-catalyzed alkene hydroalkylation.

Xi Lu received his Ph.D. degree from the University of Science and Technology of China in 2016. He is currently an Associate Researcher at the University of Science and Technology of China. His research mainly focuses on base-metal-catalyzed alkyl-alkyl bond formation.

Yao Fu received his Ph.D. degree from the University of Science and Technology of China in 2005. He is currently a Professor at the University of Science and Technology of China. His research focuses on physical organic chemistry, green organic synthesis, and the biomass chemical industry.

References

- [1] Roughley S D, Jordan A M. The medicinal chemist's toolbox: An analysis of reactions used in the pursuit of drug candidates. *J. Med. Chem.*, **2011**, *54* (10): 3451–3479.
- [2] Panfilov A V, Markovich Y D, Ivashov I P, et al. Sodium borohydride in reductive amination reactions. *Pharm. Chem. J.*, **2000**, *34* (2): 76–78.
- [3] Salvatore R N, Yoon C H, Jung K W. Synthesis of secondary amines. *Tetrahedron*, **2001**, *57* (37): 7785–7811.
- [4] Guillena G, Ramón D J, Yus M. Hydrogen autotransfer in the N-alkylation of amines and related compounds using alcohols and amines as electrophiles. *Chem. Rev.*, **2010**, *110* (3): 1611–1641.
- [5] Mao R, Frey A, Balon J, et al. Decarboxylative C(sp³)-N cross-coupling via synergetic photoredox and copper catalysis. *Nat. Catal.*, **2018**, *1* (2): 120–126.
- [6] Surry D S, Buchwald S L. Biaryl phosphane ligands in palladium-catalyzed amination. *Angew. Chem. Int. Ed.*, **2008**, *47* (34): 6338–6361.
- [7] Hartwig J F. Evolution of a fourth generation catalyst for the amination and thioetherification of aryl halides. *Acc. Chem. Res.*, **2008**, *41* (11): 1534–1544.
- [8] Surry D S, Buchwald S L. Dialkylbiaryl phosphines in Pd-catalyzed amination: A user's guide. *Chem. Sci.*, **2011**, *2* (1): 27–50.
- [9] Ley S V, Thomas A W. Modern synthetic methods for copper-mediated C(aryl)-O, C(aryl)-N, and C(aryl)-S bond formation. *Angew. Chem. Int. Ed.*, **2003**, *42* (44): 5400–5449.
- [10] Surry D S, Buchwald S L. Diamine ligands in copper-catalyzed reactions. *Chem. Sci.*, **2010**, *1* (1): 13–31.
- [11] Zhou W, Fan M, Yin J et al. CuI/oxalic diamide catalyzed coupling reaction of (hetero)aryl chlorides and amines. *J. Am. Chem. Soc.*, **2015**, *137* (37): 11942–11945.
- [12] Fan M, Zhou W, Jiang Y, et al. Assembly of primary (hetero)arylamines via CuI/oxalic diamide-catalyzed coupling of aryl chlorides and ammonia. *Org. Lett.*, **2015**, *17* (23): 5934–5937.
- [13] Yang C T, Fu Y, Huang Y B et al. Room-temperature copper-catalyzed carbon–nitrogen coupling of aryl iodides and bromides promoted by organic ionic bases. *Angew. Chem. Int. Ed.*, **2009**, *48* (40): 7398–7401.
- [14] Yu H Z, Jiang Y Y, Fu Y et al. Alternative mechanistic explanation for ligand-dependent selectivities in copper-catalyzed N- and O-arylation reactions. *J. Am. Chem. Soc.*, **2010**, *132* (51): 18078–18091.
- [15] Cheung C W, Hu X. Amine synthesis via iron-catalysed reductive coupling of nitroarenes with alkyl halides. *Nat. Commun.*, **2016**, *7*: 12494.
- [16] Cheung C W, Ploeger M, Hu X. Direct amidation of esters with nitroarenes. *Nat. Commun.*, **2017**, *8*: 14878.
- [17] Yadav M R, Nagaoka M, Kashiwara M et al. The Suzuki–Miyaura coupling of nitroarenes. *J. Am. Chem. Soc.*, **2017**, *139* (28): 9423–9426.
- [18] Rauser M, Ascheberg C, Niggemann M. Electrophilic amination with nitroarenes. *Angew. Chem. Int. Ed.*, **2017**, *56* (38): 11570–11574.
- [19] Xuan M, Lu C, Lin B L. C-S coupling with nitro group as leaving group via simple inorganic salt catalysis. *Chin. Chem. Lett.*, **2020**, *31* (1): 84–90.
- [20] Wang B, Ma J, Ren H, et al. Chemo-, site-selective reduction of nitroarenes under blue-light, catalyst-free conditions. *Chin. Chem. Lett.*, **2022**, *33* (5): 2420–2424.
- [21] Shevick S L, Wilson C V, Kotesova S, et al. Catalytic hydrogen atom transfer to alkenes: a roadmap for metal hydrides and radicals. *Chem. Sci.*, **2020**, *11* (46): 12401–12422.
- [22] Liu R Y, Buchwald S L. CuH-catalyzed olefin functionalization: From hydroamination to carbonyl addition. *Acc. Chem. Res.*, **2020**, *53* (6): 1229–1243.
- [23] Zhang Z, Bera S, Fan C, et al. Streamlined alkylation via nickel-hydride-catalyzed hydrocarbonation of alkenes. *J. Am. Chem. Soc.*, **2022**, *144* (16): 7015–7029.
- [24] He Y, Chen J, Jiang X, et al. Enantioselective NiH-catalyzed reductive hydrofunctionalization of alkenes. *Chin. J. Chem.*, **2022**, *40* (5): 651–661.
- [25] Sun C, Yin G. Integrating aryl chlorides into nickel-catalyzed 1, 1-difunctionalization of alkenes. *Chin. Chem. Lett.*, **2022**, *33* (12): 5096–5100.
- [26] Ni S X, Li Y L, Ni H Q, et al. Nickel-catalyzed hydromonofluoromethylation of unactivated alkenes for expedient construction of primary alkyl fluorides. *Chin. Chem. Lett.*, **2022** (In press). <https://doi.org/10.1016/j.ccl.2022.06.037>.
- [27] Ren X, Liu Q, Wang Z, et al. Visible-light-induced direct hydrodifluoromethylation of alkenes with difluoromethyltriphenylphosphonium iodide salt. *Chin. Chem. Lett.*, **2023**, *34* (1): 107473.
- [28] Yu Y, Jiang Y, Wu S, et al. Electrochemistry enabled selective vicinal fluorosulfonylation and fluorosulfoxidation of alkenes. *Chin. Chem. Lett.*, **2022**, *33* (4): 2009–2014.
- [29] Huang L, Arndt M, Gooßen K, et al. Late transition metal-catalyzed hydroamination and hydroamidation. *Chem. Rev.*, **2015**, *115* (7): 2596–2697.
- [30] Pirnot M T, Wang Y M, Buchwald S L. Copper hydride catalyzed hydroamination of alkenes and alkynes. *Angew. Chem. Int. Ed.*, **2016**, *55* (1): 48–57.
- [31] Kawatsura M, Hartwig J F. Palladium-catalyzed intermolecular hydroamination of vinylarenes using arylamines. *J. Am. Chem. Soc.*, **2000**, *122* (39): 9546–9547.
- [32] Waser J, Gaspar B, Nambu H, et al. Hydrazines and azides via the metal-catalyzed hydrohydrazination and hydroazidation of olefins. *J. Am. Chem. Soc.*, **2006**, *128* (35): 11693–11712.
- [33] Yang Y, Shi S L, Niu D, et al. Catalytic asymmetric hydroamination of unactivated internal olefins to aliphatic amines. *Science*, **2015**, *349*: 62–66.
- [34] Xi Y, Ma S, Hartwig J F. Catalytic asymmetric addition of an amine N–H bond across internal alkenes. *Nature*, **2020**, *588* (7837): 254–260.
- [35] Musacchio A J, Lainhart B C, Zhang X, et al. Catalytic intermolecular hydroaminations of unactivated olefins with secondary alkyl amines. *Science*, **2017**, *355* (6326): 727–730.
- [36] Gui J, Pan C M, Jin Y, et al. Practical olefin hydroamination with nitroarenes. *Science*, **2015**, *348* (6237): 886–891.
- [37] Xiao J, He Y, Ye F, et al. Remote sp³ C–H amination of alkenes with nitroarenes. *Chem*, **2018**, *4* (7): 1645–1657.
- [38] Meng L, Yang J, Duan M, et al. Facile synthesis of chiral arylamines, alkylamines and amides by enantioselective NiH-catalyzed hydroamination. *Angew. Chem. Int. Ed.*, **2021**, *60* (44): 23584–23589.
- [39] Zhang Y, Qiao D, Duan M, et al. Enantioselective synthesis of α -aminoboronates by NiH-catalysed asymmetric hydroamidation of alkenyl boronates. *Nat. Commun.*, **2022**, *13*: 5630.
- [40] Wang X X, Lu X, Li Y, et al. Recent advances in nickel-catalyzed reductive hydroalkylation and hydroarylation of electronically unbiased alkenes. *Sci China Chem*, **2020**, *63* (11): 1586–1600.
- [41] Lu X, Xiao B, Shang R, et al. Synthesis of unnatural amino acids

- through palladium-catalyzed C(sp³)-H functionalization. *Chin. Chem. Lett.*, **2016**, 27 (3): 305–311.
- [42] Lu X, He S J, Cheng W M, et al. Transition-metal-catalyzed C–H functionalization for late-stage modification of peptides and proteins. *Chin. Chem. Lett.*, **2018**, 29 (7): 1001–1008.
- [43] Chang Z, Wang J, Lu X, et al. Synthesis of *gem*-difluoroalkenes through nickel-promoted electrochemical reductive cross-coupling. *Chin. J. Org. Chem.*, **2022**, 42 (1): 147–159.
- [44] He S J, Pi J J, Li Y et al. Nickel-catalyzed Suzuki-type cross coupling of fluorinated alkenyl boronates with alkyl halides. *Acta Chim. Sinica*, **2018**, 76 (12): 956–961.
- [45] Lu X, Xiao B, Zhang Z, et al. Practical carbon–carbon bond formation from olefins through nickel-catalyzed reductive olefin hydrocarbonation. *Nat. Commun.*, **2016**, 7: 11129.
- [46] He S J, Wang J W, Li Y, et al. Nickel-catalyzed enantioconvergent reductive hydroalkylation of olefins with α -heteroatom phosphorus or sulfur alkyl electrophiles. *J. Am. Chem. Soc.*, **2020**, 142 (1): 214–221.
- [47] Wang X X, Yu L, Lu X, et al. NiH-catalyzed reductive hydrocarbonation of enol esters and ethers. *CCS Chem.*, **2022**, 4 (2): 605–615.
- [48] Wang J W, Li Y, Nie W, et al. Catalytic asymmetric reductive hydroalkylation of enamides and enecarbamates to chiral aliphatic amines. *Nat. Commun.*, **2021**, 12: 1313.
- [49] Wang J W, Liu D G, Chang Z, et al. Nickel-catalyzed switchable site-selective alkene hydroalkylation by temperature regulation. *Angew. Chem. Int. Ed.*, **2022**, 61 (31): e202205537.
- [50] Wang X X, Xu Y T, Zhang Z L, et al. NiH-catalysed proximal-selective hydroalkylation of unactivated alkenes and the ligand effects on regioselectivity. *Nat. Commun.*, **2022**, 13: 1890.
- [51] Li Y, Liu D G, Wan L, et al. Ligand-controlled cobalt-catalyzed regiodivergent alkyne hydroalkylation. *J. Am. Chem. Soc.*, **2022**, 144 (30): 13961–13972.
- [52] Wei D, Yang P, Yu C, et al. *N*-alkylation of amines with alcohols catalyzed by manganese(II) chloride or bromopentacarbonylmanganese(I). *J. Org. Chem.*, **2021**, 86 (3): 2254–2263.
- [53] Zou Q, Liu F, Zhao T, et al. Reductive amination of ketones/aldehydes with amines using BH₃N(C₂H₅)₃ as a reductant. *Chem. Commun.*, **2021**, 57 (69): 8588–8591.
- [54] Hao X W, Yuan J, Yu G A, et al. Air-stable and highly efficient indenyl-derived phosphine ligand: Application to Buchwald–Hartwig amination reactions. *J. Organomet. Chem.*, **2012**, 706–707: 99–105.